



**University of  
Zurich**<sup>UZH</sup>

**Zurich Open Repository and  
Archive**

University of Zurich  
University Library  
Strickhofstrasse 39  
CH-8057 Zurich  
[www.zora.uzh.ch](http://www.zora.uzh.ch)

---

Year: 2015

---

## **Effects of body region and time on hair cortisol concentrations in chimpanzees (*Pan troglodytes*)**

Carlitz, Esther H D ; Kirschbaum, Clemens ; Miller, Robert ; Rukundo, Joshua ; van Schaik, C P

**Abstract:** Hair cortisol concentrations (HCC) are increasingly recognized as an integrated measure of the systemic cortisol secretion. Yet, we still know very little about confounding effects on HCC in animals. The present study therefore used hair from semi-wild and zoo living chimpanzees to investigate (1) intra-individual variability of HCC (body-region effect), and (2) the stability of HCC along the hair shaft (traditionally called the washout effect). Our results indicate that absolute HCC varied substantially between certain body regions, but a factor analysis revealed that these HCC differences were mainly attributable to one common source of variance. Thus, hair from all body regions provides similar biological signals and can be mixed, albeit at the cost of a lower signal-to-noise ratio. With regard to potential underlying mechanisms, we studied skin blood flow, as observed through thermal images from one chimpanzee. We found the general HCC pattern was reflected in differences in surface body temperature observed in this individual in three out of four body regions. In a separate set of samples, we found first evidence to suggest that the systematic cortisol decrease along the hair shaft, as observed in humans, is also present in chimpanzee hair. The effect was more pronounced in semi-wild than in zoo chimpanzees presumably due to more exposure to ambient weather conditions.

DOI: <https://doi.org/10.1016/j.ygcen.2015.09.022>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-113368>

Journal Article

Accepted Version

Originally published at:

Carlitz, Esther H D; Kirschbaum, Clemens; Miller, Robert; Rukundo, Joshua; van Schaik, C P (2015). Effects of body region and time on hair cortisol concentrations in chimpanzees (*Pan troglodytes*). *General and Comparative Endocrinology*, 223:9-15.

DOI: <https://doi.org/10.1016/j.ygcen.2015.09.022>

## Accepted Manuscript

Effects of body region and time on hair cortisol concentrations in chimpanzees  
(*Pan troglodytes*)

Esther H.D. Carlitz, Clemens Kirschbaum, Robert Miller, Joshua Rukundo,  
Carel P. van Schaik


PII: S0016-6480(15)00266-X  
DOI: <http://dx.doi.org/10.1016/j.ygcen.2015.09.022>  
Reference: YGCEN 12202

To appear in: *General and Comparative Endocrinology*

Received Date: 18 May 2015  
Revised Date: 7 September 2015  
Accepted Date: 22 September 2015

Please cite this article as: Carlitz, E.H.D., Kirschbaum, C., Miller, R., Rukundo, J., van Schaik, C.P., Effects of body region and time on hair cortisol concentrations in chimpanzees (*Pan troglodytes*), *General and Comparative Endocrinology* (2015), doi: <http://dx.doi.org/10.1016/j.ygcen.2015.09.022>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



**Effects of body region and time on hair cortisol concentrations in  
chimpanzees (*Pan troglodytes*)**

**Esther H. D. Carlitz<sup>1, 2</sup>, Clemens Kirschbaum<sup>2</sup>, Robert Miller<sup>2</sup>, Joshua Rukundo<sup>3</sup>,  
Carel P. van Schaik<sup>1</sup>**

<sup>1</sup>Anthropological Institute and Museum, University of Zurich, Switzerland

<sup>2</sup>Department of Psychology, Technische Universität Dresden, Germany

<sup>3</sup>Chimpanzee Sanctuary and Wildlife Conservation Trust, Entebbe, Uganda

---

**1 Corresponding Author**

2 Esther H.D. Carlitz

3 Anthropological Institute and Museum

4 University of Zurich

5 Winterthurerstrasse 190

6 CH-8057 Zurich, Switzerland

7 Email: [ecarlitz@janegoodall.ch](mailto:ecarlitz@janegoodall.ch)

8 Phone: +41 44 635 5411 Fax: +41 44 635 6894

9

10 **Abstract:**

11 Hair cortisol concentrations (HCC) are increasingly recognized as an integrated  
12 measure of the systemic cortisol secretion. Yet, we still know very little about  
13 confounding effects on HCC in animals. The present study therefore used hair from  
14 semi-wild and zoo living chimpanzees to investigate (1) intra-individual variability of  
15 HCC (body-region effect)<sub>1</sub> and (2) the stability of HCC along the hair shaft  
16 (traditionally called the washout effect). Our results indicate that absolute HCC varied  
17 substantially between certain body regions, but a factor analysis revealed that these  
18 HCC differences were mainly attributable to one common source of variance. Thus,  
19 hair from all body regions provides similar biological signals and can be mixed, albeit  
20 at the cost of a lower signal-to-noise ratio. With regard to potential underlying  
21 mechanisms, we studied skin blood flow, as observed through thermal images from  
22 one chimpanzee. We found the general HCC pattern was reflected in differences in  
23 surface body temperature observed in this individual in three out of four body regions.  
24 In a separate set of samples, we found first evidence to suggest that the systematic  
25 cortisol decrease along the hair shaft, as observed in humans, is also present in  
26 chimpanzee hair. The effect was more pronounced in semi-wild than in zoo  
27 chimpanzees presumably due to more exposure to ambient weather conditions.

28 **Keywords**

29 Hair cortisol, stress, washout, body region

30

## 1 Introduction

Cortisol concentration measured in hair is increasingly recognized as a measure of long-term stress. Recently, the repetitive ACTH application in cows, chipmunks and lynx resulted in significantly increased HCC has provided direct evidence that hair cortisol concentrations (HCC) reflect the integrated activity of the hypothalamic-pituitary-adrenal axis (del Rosario Gonzalez-de-la-Vara et al., 2011; Mastromonaco et al., 2014; Terwissen et al., 2013). Hair cortisol analysis is increasingly applied in various circumstances, including clinical diagnosis in livestock (Comin et al., 2013), behavioral science (Dettmer et al., 2014), captive management (e.g., Carlitz et al., 2014), and conservation (e.g., Bryan et al., 2013). Hair cortisol analysis has also been applied in a growing number of mammalian species, including humans (review: Staufenbiel et al., 2013), non-human primates (e.g., Carlitz et al., 2014; Davenport et al., 2008, 2006; Dettmer et al., 2014; Fourie and Bernstein, 2011; Yamanashi et al., 2013), bears (e.g., Bechshøft et al., 2012, 2011; Macbeth et al., 2012, 2010; Malcolm et al., 2013), cows (Moya et al., 2013), horses (Anielski, 2008), dogs (Accorsi et al., 2008; Bennett and Hayssen, 2010), rock hyraxes (Koren et al., 2008), lynx (Terwissen et al., 2013) and chipmunks (Martin and Réale, 2008; Mastromonaco et al., 2014).

Despite the increased application of HCC analysis, the indicator value of HCC would be greatly improved if we knew which other factors than long-term stress affect the incorporation of and retention of cortisol in hair. Unfortunately, we insufficiently understand the factors that may confound HCC values. Various factors have been suggested, such as a systematic decline along the hair shaft (the so-called washout effect), systematic variation among body regions (body-region effect), or external contamination through sweat.

The present study attempts to fill those methodological gaps of HCC assessment in chimpanzees. While the influence of sweat is debated for humans (Grass et al., 2015; Russell et al., 2013) and horses (Anielski, 2008), it is rather unlikely to affect great ape species since their active sweat glands are mainly restricted to their palms, forehead and auxiliary organs (Montagna, 1972). We therefore examined in chimpanzees whether HCC show systematic variation across different body regions and whether HCC decreases along the hair shaft.

The washout effect has been observed in several human studies (Dettenborn et al., 2010; Gao et al., 2010; Kirschbaum et al., 2009; Skoluda et al., 2012; Steudte et al., 2011, but see Manenschijn et al., 2011; Thomson et al., 2010). It was initially suggested that structural damage of distal hair segments might permit liquids like water and cosmetics (Kirschbaum et al., 2009), or alcohol used to clean the samples before analysis (Manenschijn et al., 2011), to penetrate and wash out cortisol more easily. This would be reflected in decreasing HCC from skin-near to distal parts of the hair. While evidence for this assumption from *in vivo* studies is still pending, *in vitro* experiments by Li and colleagues (2012) showed that hair treated with shampoo solution, warm or hot water only, as well as UV-irradiation resulted in significant HCC loss, suggesting that cortisol may not only leach from the hair shaft but that it may also be degraded through UV-irradiation as was shown earlier for cannabinoids in hair (Skopp et al., 2000). Therefore, the term “waning effect” may be more appropriate than the traditionally used term “washout effect”, which was meant to indicate the leaching of cortisol by water (Kirschbaum et al., 2009).

Concerning the waning effect in animal hair, an HCC decrease by water was shown only in one *in vitro* experiment where human-like extensive wash-dry procedures resulted in decreased HCC in hair from rhesus macaques (Hamel et al., 2011). All *in*

*vivo* animal studies showed no HCC difference between the proximal and distal end of hair (Bennett and Hayssen, 2010; Carlitz et al., 2014; Davenport et al., 2006; Yamanashi et al., 2013). The latter findings may be in line with the *in vitro* experiments on hair of humans and rhesus macaques because the captive subjects were not exposed to rain and presumably not to strong UV-irradiation either. A systematic HCC decrease along the hair shaft should only be expected if hair is frequently exposed to water (Hamel et al., 2011), or UV-light (Li et al., 2012). However, Macbeth et al. (2010) also found no systematic decline along the hair shafts of water and light exposed free ranging grizzly bears, which raises the question whether some animal hair is inert to systematic HCC decrease along the hair shaft. The present study therefore investigated whether systematic cortisol decrease existed in chimpanzee hair and, if present, whether the effect could be explained by exposure to water only.

Concerning the body-region effect, an increasing number of studies suggest that HCC differs among various body regions in animals (Macbeth et al., 2010; Moya et al., 2013; Terwissen et al., 2013; Yamanashi et al., 2013, but see Carlitz et al., 2014; Comin et al., 2012; Macbeth et al., 2012), or between head regions in humans (Li et al., 2012). However, to our knowledge it has never been investigated whether HCC from all body regions provide similar biological information and what underlying mechanism may lead to this body-region effect. Yet, a better causal understanding of the effect may improve our understanding of the cortisol incorporation mechanisms into hair, which is still poorly understood (review: Meyer and Novak, 2012; Russell et al., 2012; Stalder and Kirschbaum, 2012).

Following the multiple compartment model of mechanisms of drug incorporation into hair (Henderson, 1993), cortisol in hair is discussed of being enriched from sweat

cortisol (Russell et al., 2013, but see Grass et al., 2015), from sebum, or from local cortisol production in the skin (Pang et al., 2014; Rousseau et al., 2007; Slominski et al., 2013, 2007) and hair follicles (Ito et al., 2005). Another mechanism suggested to contribute to the incorporation of cortisol into hair is the passive diffusion from the supplying blood capillaries into the growing hair cells (reviewed in: Meyer and Novak, 2012; Russell et al., 2012; Stalder and Kirschbaum, 2012), which would be in line with the fact that multiple weekly ACTH injections resulted in increased HCC in the hair segments representing the time of application (del Rosario Gonzalez-de-la-Vara et al., 2011; Mastromonaco et al., 2014; Terwissen et al., 2013). Following this idea, a higher activity of the hypothalamic-pituitary-adrenal axis leads to higher cortisol concentrations in the blood stream, which results in higher cortisol incorporation into the hair shaft due to increased diffusion pressure. In parallel, it is conceivable that differences in skin blood flow (SkBF) can influence the cortisol uptake into hair in a similar process. Increased SkBF may increase cortisol availability per time unit, which could result in higher diffusion rates and thus higher cortisol incorporation into the growing hair. Several studies on humans and various animal species have documented differences in SkBF between body regions using laser Doppler velocimetry or photopulse plethysmography (Monteiro-Riviere et al., 1990; Tsuchida, 1987; Tur et al., 1983). Because higher SkBF leads to increased heat dissipation, differences in SkBF can be indirectly measured as skin temperature (Rubinstein and Sessler, 1990). In order to test this skin blood flow hypothesis, the present study compared the HCC in four different body regions with their skin temperatures in one female chimpanzee.

We obtained hair cortisol measures from European zoo chimpanzees as well as from semi-wild living sanctuary chimpanzees (Ngamba Island, Uganda) from various body



regions in two consecutive years. Longitudinal segmentation of hair samples allowed investigating HCC measures along the hair shaft. Because the semi-wild chimpanzees were far more exposed to sun and rain than the zoo-living conspecifics, we could also examine whether this difference affected the waning effect.

## 2 Methods

### 2.1 Hair sampling and animals

Hair samples were cut from semi-wild living chimpanzees from the Ngamba Island (NI) sanctuary (Uganda) during anesthesia for the annual routine health checks in late February to early March 2011 (18 males aged 8-28 years, 20 females aged 5 to 28 years) and in late March 2012 (1 male aged 20, 9 females aged 9 to 25 years). These animals live in a 40 ha forest during the day and are encouraged with food to sleep in the roofed holding facilities at night. Hair samples were cut from six different body regions per individual (right and left dorsal forearm, center of back, right and left shoulder blade, middle of chest; there were 9 missing samples: right or left shoulder blade in 5 animals, right forearm in 1 animal, back in 1 animal, back and chest in 1 animal). For investigations on HCC stability along the hair shaft, additional hair samples were collected from 78 chimpanzees from 19 European zoos. However, only 24 samples from 13 zoos provided sufficient length and amount of material to be included in the segmentation study. Those samples were cut from the shoulder region (3 males aged 16 and 33 years, 6 females aged 3 to 43 years), back (3 males aged 30 and 41 years, 3 females aged 14 to 17 years), forearm (5 males aged 10 and 43 years) or combed from across several regions (3 males aged 17 and 36 years, 1 females aged 17 years) either in cooperation with well-trained animals ( $n = 16$ ) or during anesthesia ( $n = 5$ ) due to medical treatment.

## 2.2 Hair cortisol analysis

For studies on body-region effect, the 3 cm of hair proximal to the skin were analyzed, whereas for investigations on HCC stability along the hair shaft, the 4 cm of hair proximal to the skin were cut into 1-cm-segments. Here, all hair strands that provided sufficient length and amount of material were included in the analysis ( $n_{\text{Zoo}} = 24$ ,  $n_{\text{NI2011}} = 46$ ,  $n_{\text{NI2012}} = 25$ ).

For cortisol analysis, all subsamples were washed twice for three minutes with 3 ml isopropanol. The air-dried hair was then minced into 3-5 mm pieces and 6 mg of this 10-50 mg pool were incubated with 1.8 ml methanol for 17 h in a glass tube. 1.6 ml of the extract were transferred into another glass tube, dried and re-suspended in 150  $\mu$ l Aqua bidest. Cortisol concentrations were determined using a commercially available immunoassay with luminescence detection (LIA, IBL-Hamburg, Germany). Intra- and inter-assay coefficients of variation of this assay are below 8%.

## 2.3 Heat dissipation measures

Thermal images from the front and the back were taken from one female zoo chimpanzee using a thermographic camera (Vario Cam® high resolution, InfraTec GmbH, Dresden, Germany). The Software IRBIS ® 3 (InfraTec) was employed to extract temperature measures from the chest, forearm, shoulder blade and back according to the hair sampling points from the Ngamba Island chimpanzees. Measure points were directed to areas with little hair, which exhibited the highest temperature in order to avoid false measures due to the insulating effect of fur. For each body region, the mean temperature with standard deviation was derived from a small circular area (supplemental material, Figure s1).

For this study, it was not possible to add more thermal images from other animals because either chimpanzees had too much fur which prevented us from obtaining

clear measures or there was no thermographic camera with high resolution available in the facilities.

## 2.4 Statistical analysis

Hair cortisol data was not found to be normally distributed. Fourth root transformation reduced the skewness of distribution most effectively and was applied prior to data analyses (Miller and Plessow, 2013). For descriptive purposes, information on mean values and standard deviations are presented in original units (pg/mg). All analyses were performed using R 3.1.1 (R Core Team, 2015) statistical software.

### 2.4.1 HCC stability along hair shaft

Mixed-effects linear regression models were employed to assess potential HCC changes across the four consecutive segments in both zoo and NI samples, while accounting for systematic variance due to different chimpanzees and body regions. Thus, the stability of HCC across the four consecutive segments could be quantified by means of intra-class correlation coefficients (ICC; see Hruschka et al., 2005). The upper bound of such ICCs (i.e. ICC = 1) indicates perfect stability of HCC across segments (i.e., the rank order of HCCs from the first segments is also preserved across the subsequent segments), whereas the lower bound (ICC = 0) indicates no association between the different segments. The employed model is expressed in equation 1:

$$HCC^{0.25}_{ijk} = \beta_0 + \beta_1 \times \text{Segment} + \xi_i + \zeta_{ij} + \varepsilon_{ijk} \quad (1)$$

where  $\beta_0$  denotes  $HCC^{0.25}$  in the first segment,  $\beta_1$  denotes the change of  $HCC^{0.25}$  per subsequent segment. The random effects  $\xi_i$  and  $\zeta_{ij}$  account for the variation in  $HCC^{0.25}$  due to differences between individuals  $i$  and body-regions  $j$ , respectively.

## 2.4.2 HCC in different body regions

For descriptive purposes, Pearson correlation coefficients were calculated to investigate the HCC relationships between the different body regions. Thereafter, a confirmatory factor analysis was performed (Rosseel, 2012) to investigate whether the covariance structure of HCC across different body regions could be sufficiently accounted for by one common factor. Besides the robust likelihood ratio statistic (Bentler and Yuan, 1999), model fit was evaluated using the comparative fit index (CFI), root mean square error of approximation (RMSEA), standardized root mean square residual (SRMR) and the Bayesian information criterion (BIC). HCC differences between the investigated body regions were assessed using paired t-Tests.

## 3 Results

### 3.1 HCC stability along the hair shaft

Pair-wise comparison revealed that all four segments were highly correlated in both samples from the zoos and from Ngamba Island ( $0.9 \geq r \geq 0.7$ , all  $p$ 's  $< 0.001$ ). Mixed-effects regressions of HCC on the four consecutive 1-cm-segments revealed profound differences between change of HCC in chimpanzee hair from European zoos and those from the NI sanctuary. In zoo chimpanzees, we found a decrease of HCC across all segments that reached statistical significance ( $\chi^2(1) = 4.22$ ,  $p = 0.040$ ) but was numerically negligible (mean  $\pm$  SD decrease of HCC per segment:  $5.7\% \pm 18.8\%$ ; 61.1% explained variance). The intra-individual stability of HCC across the four segments amounted to ICC = 0.81. Hair samples from NI-chimpanzees showed a more pronounced cortisol decline along the hair strand ( $\chi^2(1) = 72.40$ ,  $p < 0.001$ ; mean  $\pm$  SD decrease of HCC per segment:  $20.1\% \pm 24.9\%$ ; 69.6% explained variance). However, the intra-individual stability of HCC across the

four segments was similar to that of the zoo samples, and amounted to  $ICC = 0.83$ . After controlling for body-region-related differences in HCC, the intra-individual stability estimate of HCC drops to  $ICC = 0.32$ . This is the proportion of HCC that can actually be attributed to environmental and individual stability across the time interval covered by the hair stands from different body regions of a single chimpanzee. The mean trajectories of the HCC decrease across segments, as traditionally reported in previous studies (Carlitz et al., 2014; Kirschbaum et al., 2009; Manenschijn et al., 2011), are visualized in Figure 1 whereas a more detailed visualization of the HCC development along segments is presented in Figure 2.

[Please insert Figure 1 here]

[Please insert Figure 2 here]

### 3.2 HCC in different body regions

Descriptive statistics for HCC in the different body regions are provided in Table 1. Pearson correlations were highly significant between all pairs of body regions (Table ) and showed a strong effect size (Cohen, 1988). Results from the factor-analysis revealed that the HCC covariance structure could be sufficiently accounted for by only one factor ( $\chi^2(2) = 2.07, p = 0.35$ ;  $CFI = 1.00$ ,  $RMSEA_{90\%} = 0.00 - 0.31$ ,  $SRMR = 0.02$ ,  $BIC = -119.74$ ), suggesting that HCC at different body regions were driven by one systemic factor across the time interval covered by the length of hair strand. Factor loadings indicated, that chest HCC ( $\lambda = 0.90, p < 0.01$ ) contributed most substantially to this “cortisol exposure” factor, followed by shoulder and forearm HCC (both  $\lambda$ 's =  $0.87, p$ 's  $< 0.01$ ), and finally back HCC ( $\lambda = 0.78, p < 0.01$ ). Constraining all loadings to equality did not result in a significantly worse model fit ( $\Delta\chi^2(3) = 6.27, p = 0.10$  with  $\chi^2(5) = 8.95, p = 0.11$ , with ;  $CFI = 0.98$ ,  $RMSEA_{90\%} = 0.00 - 0.25$ ;  $SRMR$

252 = 0.13, BIC = -122.56). Therefore, HCC differences between animals probably are  
253 equally manifested in hair from all body regions.

254 [Please insert Table 1 here.]

255 Besides the strong correlations, the paired t-Tests revealed that the absolute HCC  
256 differed (at least marginally) significantly between all investigated regions (all  $p$ 's  $\leq$   
257 0.06; Figure 3). HCC was significantly higher in the chest region followed by  
258 shoulder, forearm and back. A very similar pattern was observed for temperature  
259 measures based on thermal images from one chimpanzee although only three out of  
260 four body regions followed the average HCC pattern in this particular chimpanzee  
261 (Figure 4, Figure s1).

262 [Please insert Figure 3 and Figure 4 here.]

## 263 **4 Discussion**

264 Hair cortisol analysis is increasingly applied to measure long-term stress in animals  
265 and humans, but we still lack sufficient understanding of potentially confounding  
266 effect, such as the waning effect along the hair shaft and possible differences in  
267 steroid incorporation at different body regions. In this study, we estimated these  
268 effects using chimpanzee hairs.

269 Concerning the waning effect, our results revealed a strong HCC decrease towards  
270 the distal end of the hair shaft in samples from Ngamba Island chimpanzees as well  
271 as a significant though very weak effect in samples from European zoo chimpanzees.  
272 While this is in line with many human studies (Dettenborn et al., 2012; Kirschbaum et  
273 al., 2009; Skoluda et al., 2012, but see Manenschijn et al., 2011; Thomson et al.,  
274 2010), the present results oppose all previous animal studies, which reported no

systematic cortisol decrease towards distal segments (Bennett and Hayssen, 2010; Carlitz et al., 2014; Davenport et al., 2006; Macbeth et al., 2010; Yamanashi et al., 2013).

It is unlikely that differences in laboratory protocols, as suggested by Manenschijn et al. (2011), led to the discrepancy between the present and former results because the same protocol was also applied in our orangutan study, in which no cortisol decrease along 15 cm long hair shafts was observed (Carlitz et al., 2014). It is also unlikely that this is a species specific result since Yamanashi et al. (2013) did not find a waning effect in chimpanzees either. There might be a difference between captive animals that are sheltered from ambient weather conditions and (semi-) wild animals that are exposed to ambient weathers. However, Macbeth et al. (2010), the only study investigating HCC along the hair shaft in wild animals, did not find a waning effect either. Yet, it is conceivable that the formerly applied statistical methods (repeated-measures ANOVA) were not sensitive enough because it pools valuable information. Interestingly, a repeated-measures ANOVA would not have brought to light the small waning effect in the present samples from zoo chimpanzees but only in the NI samples (data not shown). The detailed graph of cortisol changes along the hair shaft (Figure ) illustrates that in zoo samples the effect is masked by considerable background noise and is only visible in the second and third segment if samples with greater initial HCC are included. This may be due to the fact that individual samples with low (initial) HCC are more likely to show an increase or no change over time rather than a decrease of HCC due to bottom effects. Furthermore, it is conceivable that small quantities of cortisol exhibit stronger bonds to the hair matrix and are less likely to be removed by external factors than larger quantities of cortisol. Macbeth et al. (2010) already mentioned the possibility that a waning effect

was not detectable in their study on wild grizzly bears because of low HCC in their samples.

In search of underlying mechanisms for the systematic cortisol decrease observed in the present study, internal factors (i.e., stress related) as well as external factors (e.g., rain, sun) have to be considered. Concerning internal factors, it seems unlikely that stress levels decreasing systematically across the time period covered by the hair segments affected the whole NI group in two consecutive years and the inhomogeneous zoo group alike, resulting in systematic cortisol decrease along the hair shaft. Concerning external factors, water (hair washes in humans or rain in animals) has often been suggested as the primary source of cortisol decrease along the hair shaft, which is why this effect was called washout effect (Kirschbaum et al., 2009). To date, however, only two *in vitro* studies (Hamel et al., 2011; Li et al., 2012), revealed a direct relationship between exposure to water and HCC decrease along the hair. Moreover, the one *in vivo* study (Dettenborn et al., 2012), which tried to correlate the number of weekly hair washes in humans with the waning effect along the hair shaft, may not have found the effect because of too many other factors that add noise to the data.

The *in vitro* experiment from Li et al. (2012) suggests that water and UV-irradiation can reduce HCC independently, although *in vivo* data for the effect of UV-irradiation on HCC is still pending. The pronounced cortisol decrease in NI samples and the very weak effect in samples from zoo animals could also be interpreted as the result of difference in exposition to UV-irradiation (moderate European vs. strong equatorial sun), difference in exposition to water (negligible in European zoos vs. routinely in NI animals), or differences of a combination of both. However, this interpretation at present remains speculative because of the higher noise level (i.e., fluctuation of the



environment) in the inhomogeneous zoo samples, and a weakened hair structure from one factor with cortisol removal, or degradation from another factor (*cf.* Manenschijn et al., 2011) can also not be excluded. Thus, the underlying mechanisms for the waning effect remain to be investigated as well as the question to what extent hair samples from the European zoo animals are affected.

Despite the presence of the waning effect our results showed a strong intra-sample stability in both groups and all segments provided similar biological information (although absolute HCC differed between segments). Thus, if animals are exposed to similar ambient conditions, the effect can be ‘controlled for’ using the same length of hair throughout the study.

Concerning the body-region effect, our results revealed that absolute HCC differed significantly between body regions, which is in line with previous studies (Macbeth et al., 2010; Moya et al., 2013; Terwissen et al., 2013; Yamanashi et al., 2013). In addition, and similar to the chimpanzee study from Yamanashi et al. (2013), our results revealed strong correlations between body regions. The present data extend previous research showing that HCC measures of all body regions were mainly driven by one common factor. Thus, HCC in all body regions appear to convey information about the same biological entity, and it is likely that this biological entity is an excellent representation of the systemic cortisol secretion as suggested by an increasing body of literature in animals (e.g., Carlitz et al., 2014; del Rosario Gonzalez-de-la-Vara et al., 2011; Malcolm et al., 2013; Mastromonaco et al., 2014; Terwissen et al., 2013) and humans (e.g., O’Brien et al., 2013; Stalder et al., 2014). Assuming no asymmetry of hair samples from different regions, controlling for body region could be omitted if a large enough sample from random body regions is

available, even though it inevitably will reduce the signal-to-noise ratio. Thus, it is possible to use shed hair samples as a mixture of various body regions.

In an attempt to explain the underlying mechanism of the effect of body region, we found that skin temperature, which is a measure of skin blood flow (Rubinstein and Sessler, 1990), increased in the same order as the mean HCC levels from the 38 NI chimpanzees (back<forearm<shoulder<chest). Even though there is an unexplained mismatch for one body region between skin temperature and HCC in the particular subject for which skin temperature was recorded, it is conceivable that higher skin blood flow indicates a greater blood supply to the hair root per unit time, which might result in a higher diffusion rate and therefore in more cortisol incorporation into the hair shaft. Indeed, HCC patterns of body regions described earlier for chimpanzees (side > back > elbow; Yamanashi et al., 2013) and for orang-utans (shoulder > stomach = back > wrist, although this pattern was not significant; Carlitz et al., 2014) also seem to follow the skin temperature gradient illustrated in the thermal images of one chimpanzee (see supplemental material Figure s1). While these results encourage further investigations of the relationship between HCC and skin blood flow, more data with temperature measures from more animals as well as from more body regions are necessary to verify this skin blood flow hypothesis. Yet, future work should also consider a potential interaction between the systemic and local cortisol production as suggested by Keckeis et al. (2012), although Pang et al. (2014) found that local cortisol production in the skin was reduced with increased systemic cortisol concentration.

In conclusion, this study presents first evidence that the waning effect, so far only reported for human hair, influences HCC along the hair shaft of chimpanzees. Nonetheless, all segments provided similar biological information. Regarding the

body-region effect, our results confirm that absolute HCC differ between body regions and extend previous research by showing that HCC in all body regions convey similar biological information, presumably the systemic cortisol secretion. In conclusion, shed hair from various unknown body regions can well be used in observational studies at the cost of a lower signal-to-noise ratio.

#### **Acknowledgement:**

This study could not have been conducted without the great support from all the animal keepers, the veterinary team, and Director Lilly Ajarova from the Ngamba Island *Chimpanzee Sanctuary & Wildlife Conservation Trust*. Furthermore, we would like to thank Jutta Heuer and Cornelia Hofman from Zoo Halle as well as Steffen Kranz from the EVH GmbH Halle for their help to take thermal images from a chimpanzee. For the provisioning of chimpanzee hair samples, additional thanks go to curators, animal keepers and veterinarians from zoos and parks in Amersfoort, Amsterdam, Augsburg, Banham, Borås, Furuvik, Givskud, Halle, Heidelberg, Jerez, Kristiansand, Leintal, Liberec, Madrid, Magdeburg, Odense, Osnabrück, La Vallée des Singes, and Tallin. Funding for this study was kindly provided by the Jane Goodall Institute Switzerland and the A.H. Schultz-Stiftung to Esther Carlitz.

# References

- Accorsi, P.A., Carloni, E., Valsecchi, P., Viggiani, R., Gamberoni, M., Tamanini, C., Seren, E., 2008. Cortisol determination in hair and faeces from domestic cats and dogs. *Gen. Comp. Endocrinol.* 155, 398–402.
- Anielski, P., 2008. Hair analysis of anabolic steroids in connection with doping control—results from horse samples. *J. Mass Spectrom.* 43, 1001–1008.
- Bechshøft, T.Ø., Riget, F.F., Sonne, C., Letcher, R.J., Muir, D.C.G., Novak, M.A., Henchey, E., Meyer, J.S., Eulaers, I., Jaspers, V.L.B., others, 2012. Measuring environmental stress in East Greenland polar bears, 1892–1927 and 1988–2009: What does hair cortisol tell us? *Environ. Int.* 45, 15–21.
- Bechshøft, T.Ø., Sonne, C., Dietz, R., Born, E.W., Novak, M.A., Henchey, E., Meyer, J.S., 2011. Cortisol levels in hair of East Greenland polar bears. *Sci. Total Environ.* 409, 831–834.
- Bennett, A., Hayssen, V., 2010. Measuring cortisol in hair and saliva from dogs: coat color and pigment differences. *Domest. Anim. Endocrinol.* 39, 171–180.
- Bentler, P.M., Yuan, K.-H., 1999. Structural Equation Modeling with Small Samples: Test Statistics. *Multivar. Behav. Res.* 34, 181–197. doi:10.1207/S15327906Mb340203
- Bryan, H.M., Darimont, C.T., Paquet, P.C., Wynne-Edwards, K.E., Smits, J.E.G., 2013. Stress and Reproductive Hormones in Grizzly Bears Reflect Nutritional Benefits and Social Consequences of a Salmon Foraging Niche. *PLoS ONE* 8, e80537. doi:10.1371/journal.pone.0080537
- Carlitz, E.H.D., Kirschbaum, C., Stalder, T., van Schaik, C.P., 2014. Hair as a long-term retrospective cortisol calendar in orang-utans (*Pongo* spp.): New perspectives for stress monitoring in captive management and conservation. *Gen. Comp. Endocrinol.* 195, 151–156. doi:10.1016/j.ygcen.2013.11.002
- Cohen, J., 1988. Statistical power analysis for the behavioral science. Hillsdale, Erlbaum.
- Comin, A., Peric, T., Corazzin, M., Veronesi, M.C., Meloni, T., Zufferli, V., Cornacchia, G., Prandi, A., 2013. Hair cortisol as a marker of hypothalamic-pituitary-adrenal axis activation in Friesian dairy cows clinically or physiologically compromised. *Livest. Sci.* 152, 36–41. doi:10.1016/j.livsci.2012.11.021
- Comin, A., Zufferli, V., Peric, T., Canavese, F., Barbetta, D., Prandi, A., 2012. Hair cortisol levels determined at different body sites in the New Zealand White rabbit. *World Rabbit Sci.* 20, 149–154.
- Davenport, M.D., Lutz, C.K., Tiefenbacher, S., Novak, M.A., Meyer, J.S., 2008. A rhesus monkey model of self-injury: effects of relocation stress on behavior and neuroendocrine function. *Biol. Psychiatry* 63, 990–996.
- Davenport, M.D., Tiefenbacher, S., Lutz, C.K., Novak, M.A., Meyer, J.S., 2006. Analysis of endogenous cortisol concentrations in the hair of rhesus macaques. *Gen. Comp. Endocrinol.* 147, 255–261. doi:10.1016/j.ygcen.2006.01.005
- del Rosario Gonzalez-de-la-Vara, M., Valdez, R.A., Lemus-Ramirez, V., Vazquez-Chagoyan, J.C., Villa-Godoy, A., Romano, M.C., 2011. Effects of adrenocorticotrophic hormone challenge and age on hair cortisol concentrations in dairy cattle. *Can. J. Vet. Res.* 75, 216–221.
- Dettenborn, L., Tietze, A., Bruckner, F., Kirschbaum, C., 2010. Higher cortisol content in hair among long-term unemployed individuals compared to controls.

- 441 Psychoneuroendocrinology 35, 1404–1409.  
442 doi:10.1016/j.psyneuen.2010.04.006
- 443 Dettenborn, L., Tietze, A., Kirschbaum, C., Stalder, T., 2012. The assessment of  
444 cortisol in human hair: associations with sociodemographic variables and  
445 potential confounders. *Stress* 15, 578–588.
- 446 Dettmer, A.M., Novak, M.A., Meyer, J.S., Suomi, S.J., 2014. Population density-  
447 dependent hair cortisol concentrations in rhesus monkeys (*Macaca mulatta*).  
448 *Psychoneuroendocrinology* 42, 59–67. doi:10.1016/j.psyneuen.2014.01.002
- 449 Fourie, N.H., Bernstein, R.M., 2011. Hair cortisol levels track phylogenetic and age  
450 related differences in hypothalamic–pituitary–adrenal (HPA) axis activity in  
451 non-human primates. *Gen. Comp. Endocrinol.* 174, 150–155.  
452 doi:10.1016/j.ygcen.2011.08.013
- 453 Gao, W., Xie, Q., Jin, J., Qiao, T., Wang, H., Chen, L., Deng, H., Lu, Z., 2010. HPLC-  
454 FLU detection of cortisol distribution in human hair. *Clin. Biochem.* 43, 677–  
455 682.
- 456 Grass, J., Kirschbaum, C., Miller, R., Gao, W., Steudte-Schmiedgen, S., Stalder, T.,  
457 2015. Sweat-inducing physiological challenges do not result in acute changes  
458 in hair cortisol concentrations. *Psychoneuroendocrinology*.
- 459 Hamel, A.F., Meyer, J.S., Henchey, E., Dettmer, A.M., Suomi, S.J., Novak, M.A.,  
460 2011. Effects of shampoo and water washing on hair cortisol concentrations.  
461 *Clin. Chim. Acta* 412, 382–385.
- 462 Henderson, G.L., 1993. Mechanisms of drug incorporation into hair. *Forensic Sci. Int.*  
463 63, 19–29. doi:10.1016/0379-0738(93)90256-A
- 464 Hruschka, D.J., Kohrt, B.A., Worthman, C.M., 2005. Estimating between- and within-  
465 individual variation in cortisol levels using multilevel models.  
466 *Psychoneuroendocrinology* 30, 698–714. doi:10.1016/j.psyneuen.2005.03.002
- 467 Ito, N., Ito, T., Kromminga, A., Bettermann, A., Takigawa, M., Kees, F., Straub, R.H.,  
468 Paus, R., 2005. Human hair follicles display a functional equivalent of the  
469 hypothalamic-pituitary-adrenal axis and synthesize cortisol. *FASEB J.* 19,  
470 1332–1334. doi:10.1096/fj.04-1968fje
- 471 Keckeis, K., Lepschy, M., Schöpper, H., Moser, L., Troxler, J., Palme, R., 2012. Hair  
472 cortisol: a parameter of chronic stress? Insights from a radiometabolism study  
473 in guinea pigs. *J. Comp. Physiol. B* 182, 985–996. doi:10.1007/s00360-012-  
474 0674-7
- 475 Kirschbaum, C., Tietze, A., Skoluda, N., Dettenborn, L., 2009. Hair as a retrospective  
476 calendar of cortisol production—increased cortisol incorporation into hair in the  
477 third trimester of pregnancy. *Psychoneuroendocrinology* 34, 32–37.
- 478 Koren, L., Mokady, O., Geffen, E., 2008. Social status and cortisol levels in singing  
479 rock hyraxes. *Horm. Behav.* 54, 212–216.
- 480 Li, J., Xie, Q., Gao, W., Xu, Y., Wang, S., Deng, H., Lu, Z., 2012. Time course of  
481 cortisol loss in hair segments under immersion in hot water. *Clin. Chim. Acta*  
482 413, 434–440. doi:10.1016/j.cca.2011.10.024
- 483 Macbeth, B.J., Cattet, M.R.L., Obbard, M.E., Middel, K., Janz, D.M., 2012. Evaluation  
484 of hair cortisol concentration as a biomarker of long-term stress in free-ranging  
485 polar bears. *Wildl. Soc. Bull.* 36, 747–758. doi:10.1002/wsb.219
- 486 Macbeth, B.J., Cattet, M.R.L., Stenhouse, G.B., Gibeau, M.L., Janz, D.M., 2010. Hair  
487 cortisol concentration as a noninvasive measure of long-term stress in free-  
488 ranging grizzly bears (*Ursus arctos*): considerations with implications for other  
489 wildlife. *Can. J. Zool.* 88, 935–949. doi:10.1139/Z10-057
- 490 Malcolm, K.D., McShea, W.J., Van Deelen, T.R., Bacon, H.J., Liu, F., Putman, S.,  
491 Zhu, X., Brown, J.L., 2013. Analyses of fecal and hair glucocorticoids to

- 492 evaluate short- and long-term stress and recovery of Asiatic black bears  
 493 (*Ursus thibetanus*) removed from bile farms in China. *Gen. Comp. Endocrinol.*  
 494 185, 97–106. doi:10.1016/j.ygcen.2013.01.014
- 495 Manenschiijn, L., Koper, J.W., Lamberts, S.W., van Rossum, E.F., 2011. Evaluation  
 496 of a method to measure long term cortisol levels. *Steroids* 76, 1032–1036.
- 497 Martin, J.G.A., Réale, D., 2008. Animal temperament and human disturbance:  
 498 implications for the response of wildlife to tourism. *Behav. Processes* 77, 66–  
 499 72.
- 500 Mastromonaco, G.F., Gunn, K., McCurdy-Adams, H., Edwards, D.B., Schulte-  
 501 Hostedde, A.I., 2014. Validation and use of hair cortisol as a measure of  
 502 chronic stress in eastern chipmunks (*Tamias striatus*). *Conserv. Physiol.* 2,  
 503 cou055.
- 504 Meyer, J.S., Novak, M.A., 2012. Minireview: hair cortisol: a novel biomarker of  
 505 hypothalamic-pituitary-adrenocortical activity. *Endocrinology* 153, 4120–4127.
- 506 Miller, R., Plessow, F., 2013. Transformation techniques for cross-sectional and  
 507 longitudinal endocrine data: Application to salivary cortisol concentrations.  
 508 *Psychoneuroendocrinology* 38, 941–946.
- 509 Montagna, W., 1972. The skin of nonhuman primates. *Am. Zool.* 12, 109–124.
- 510 Monteiro-Riviere, N.A., Bristol, D.G., Manning, T.O., Rogers, R.A., Riviere, J.E.,  
 511 1990. Interspecies and interregional analysis of the comparative histologic  
 512 thickness and laser Doppler blood flow measurements at five cutaneous sites  
 513 in nine species. *J. Invest. Dermatol.* 95, 582–586.
- 514 Moya, D., Schwartzkopf-Genswein, K.S., Veira, D.M., 2013. Standardization of a  
 515 non-invasive methodology to measure cortisol in hair of beef cattle. *Livest. Sci.*  
 516 158, 138–144. doi:10.1016/j.livsci.2013.10.007
- 517 O'Brien, K.M., Tronick, E.Z., Moore, C.L., 2013. Relationship between Hair Cortisol  
 518 and Perceived Chronic Stress in a Diverse Sample. *Stress Health* 29, 337–  
 519 344. doi:10.1002/smi.2475
- 520 Pang, S., Wu, H., Wang, Q., Cai, M., Shi, W., Shang, J., 2014. Chronic Stress  
 521 Suppresses the Expression of Cutaneous Hypothalamic–Pituitary–  
 522 Adrenocortical Axis Elements and Melanogenesis.
- 523 R Core Team, 2015. R: A Language and Environment for Statistical Computing. R  
 524 Foundation for Statistical Computing, Vienna [WWW Document]. URL  
 525 <http://www.r-project.org/> (accessed 2.26.15).
- 526 Rosseel, Y., 2012. lavaan: An R package for structural equation modeling. *J. Stat.*  
 527 *Softw.* 48, 1–36.
- 528 Rousseau, K., Kauser, S., Pritchard, L.E., Warhurst, A., Oliver, R.L., Slominski, A.,  
 529 Wei, E.T., Thody, A.J., Tobin, D.J., White, A., 2007. Proopiomelanocortin  
 530 (POMC), the ACTH/melanocortin precursor, is secreted by human epidermal  
 531 keratinocytes and melanocytes and stimulates melanogenesis. *FASEB J.* 21,  
 532 1844–1856.
- 533 Rubinstein, E.H., Sessler, D.I., 1990. Skin-surface temperature gradients correlate  
 534 with fingertip blood flow in humans. *Anesthesiology* 73, 541–545.
- 535 Russell, E., Koren, G., Rieder, M., Van Uum, S., 2012. Hair cortisol as a biological  
 536 marker of chronic stress: Current status, future directions and unanswered  
 537 questions. *Psychoneuroendocrinology* 37, 589–601.  
 538 doi:10.1016/j.psyneuen.2011.09.009
- 539 Russell, E., Koren, G., Rieder, M., Van Uum, S.H.M., 2013. The Detection of Cortisol  
 540 in Human Sweat: Implications for Measurement of Cortisol in Hair. *Ther. Drug*  
 541 *Monit.* 1. doi:10.1097/FTD.0b013e31829daa0a



- Skoluda, N., Dettenborn, L., Stalder, T., Kirschbaum, C., 2012. Elevated hair cortisol concentrations in endurance athletes. *Psychoneuroendocrinology* 37, 611–617.
- Skopp, G., Pötsch, L., Mauden, M., 2000. Stability of cannabinoids in hair samples exposed to sunlight. *Clin. Chem.* 46, 1846–1848.
- Slominski, A., Wortsman, J., Tuckey, R.C., Paus, R., 2007. Differential expression of HPA axis homolog in the skin. *Mol. Cell. Endocrinol.* 265, 143–149.
- Slominski, A., Zbytek, B., Nikolakis, G., Manna, P.R., Skobowiat, C., Zmijewski, M., Li, W., Janjetovic, Z., Postlethwaite, A., Zouboulis, C.C., others, 2013. Steroidogenesis in the skin: implications for local immune functions. *J. Steroid Biochem. Mol. Biol.* 137, 107–123.
- Stalder, T., Kirschbaum, C., 2012. Analysis of cortisol in hair – State of the art and future directions. *Brain Behav Immun* 26, 1019–1029. doi:10.1016/j.bbi.2012.02.002
- Stalder, T., Tietze, A., Steudte, S., Alexander, N., Dettenborn, L., Kirschbaum, C., 2014. Elevated hair cortisol levels in chronically stressed dementia caregivers. *Psychoneuroendocrinology* 47, 26–30. doi:10.1016/j.psyneuen.2014.04.021
- Staufenbiel, S.M., Penninx, B.W.J.H., Spijker, A.T., Elzinga, B.M., van Rossum, E.F.C., 2013. Hair cortisol, stress exposure, and mental health in humans: A systematic review. *Psychoneuroendocrinology* 38, 1220–1235. doi:10.1016/j.psyneuen.2012.11.015
- Steudte, S., Stalder, T., Dettenborn, L., Klumbies, E., Foley, P., Beesdo-Baum, K., Kirschbaum, C., 2011. Decreased hair cortisol concentrations in generalised anxiety disorder. *Psychiatry Res.* 186, 310–314. doi:10.1016/j.psychres.2010.09.002
- Terwissen, C.V., Mastromonaco, G.F., Murray, D.L., 2013. Influence of adrenocorticotrophin hormone challenge and external factors (age, sex, and body region) on hair cortisol concentration in Canada lynx (*Lynx canadensis*). *Gen. Comp. Endocrinol.* 194, 162–167. doi:10.1016/j.ygcen.2013.09.010
- Thomson, S., Koren, G., Fraser, L.-A., Rieder, M., Friedman, T.C., Van Uum, S.H.M., 2010. Hair Analysis Provides a Historical Record of Cortisol Levels in Cushing's Syndrome. *Exp. Clin. Endocrinol. Diabetes Off. J. Ger. Soc. Endocrinol. Ger. Diabetes Assoc.* 118, 133–138. doi:10.1055/s-0029-1220771
- Tsuchida, Y., 1987. Regional differences in the skin blood flow at various sites of the body studied by xenon 133. *Plast. Reconstr. Surg.* 80, 705–708.
- Tur, E., Tur, M., Maibach, H.I., Guy, R.H., 1983. Basal Perfusion of the Cutaneous Microcirculation: Measurements as a Function of Anatomic Position. *J. Invest. Dermatol.* 81, 442–446. doi:10.1111/1523-1747.ep12522619
- Yamanashi, Y., Morimura, N., Mori, Y., Hayashi, M., Suzuki, J., 2013. Cortisol analysis of hair of captive chimpanzees (*Pan troglodytes*). *Gen. Comp. Endocrinol.* 194, 55–63. doi:10.1016/j.ygcen.2013.08.013




Figure 1: Mean hair cortisol concentrations ( $\pm$  SE) of the four proximal 1-cm-segments from 24 European zoos-living chimpanzees (gray line,  $n = 24$  samples), 38 chimpanzees from Ngamba Island (Uganda) in 2011 (dashed black line,  $n = 46$  samples), and from 7 Ngamba Island chimpanzees in 2012 (solid black line,  $n = 25$  samples).






Figure 2: Detailed illustration of the hair cortisol changes along four consecutive 1-cm-segments for samples from European zoo chimpanzees ( $n = 24$ ) and from semi-wild living chimpanzees from the Ngamba Island (NI) Chimpanzee Sanctuary, Uganda (hair samples from 2011,  $n = 46$ ; hair samples from 2012,  $n = 25$ ). Negative y-values indicate that cortisol concentrations (HCC) in segment 2 (a), segment 3 (b) or segment 4 (c) was smaller than cortisol concentration in segment 1. Cortisol decrease towards the distal end of hair was present in both NI ( $\chi^2(1) = 72.40$ ,  $p < 0.001$ ) and zoo samples ( $\chi^2(1) = 4.22$ ,  $p = 0.040$ ), but only NI samples exhibited significant correlations between HCC in segment 1 and the loss of HCC across all following segments. \*\*\* $p < 0.001$




Figure 3: Bar plot illustrating the mean hair cortisol concentrations ( $\pm$ SE) of four body regions from 48 samples sets from semi-wild ranging chimpanzees from the Ngamba Island chimpanzee sanctuary (Uganda; 2011:  $n = 38$ , 2012:  $n = 10$ ). Each value is an integrated measure of the three cm of hair closest to the skin. Forearm and shoulder show the mean concentrations of the right and left side. +  $p = 0.06$ , \*  $p < 0.05$ , \*\*  $p < 0.01$ , \*\*\*  $p < 0.001$ .




Figure 4: Bar plot illustrating the hair cortisol concentration (gray bars) and skin temperature ( $\pm$  SD, white bars) of four body regions from one female zoo chimpanzee. Each cortisol value is an integrated measure of the three cm of hair closest to the skin. Temperature measures were measured using thermal imaging. Forearm and shoulder samples were taken from the right side.

Table 1: Pearson correlation coefficients of hair cortisol concentrations of four body regions from 48 chimpanzees. Point estimates are listed below the main diagonal of the matrix, whereas their bootstrapped 95% confidence regions are listed above the main diagonal.

	back	forearm	shoulder	chest
N	46	48	47	47
Mean	3.41	3.88	4.92	6.88
SD	1.73	2.69	3.48	5.31
Minimum	0.18	0.6	0.97	1.09
Maximum	9.28	16.82	17.01	31.49
Correlation (back)	1	[0.62, 0.82]	[0.53, 0.77]	[0.56, 0.80]
Correlation (forearm)	0.73	1	[0.57, 0.87]	[0.60, 0.88]
Correlation (shoulder)	0.67	0.75	1	[0.68, 0.89]
Correlation (chest)	0.69	0.78	0.81	1

620 **Highlights:**

621 Cortisol concentrations decreased significantly along the hair shaft of chimpanzees.

622 All segments along the hair shaft provided similar biological information.

623 Absolute hair cortisol concentrations (HCC) differed between body regions.

624 HCC among various body regions conveyed similar biological information.

625

626

627